



Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena, A. Provenzale

► To cite this version:

M. Baudena, A. Provenzale. Rainfall intermittency and vegetation feedbacks in drylands. Hydrology and Earth System Sciences Discussions, 2007, 4 (6), pp.4241-4264. hal-00298914

HAL Id: hal-00298914

<https://hal.science/hal-00298914>

Submitted on 22 Nov 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Papers published in *Hydrology and Earth System Sciences Discussions* are under open-access review for the journal *Hydrology and Earth System Sciences*

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena¹ and A. Provenzale²

¹CIMA, Centro di ricerca Interuniversitario in Monitoraggio Ambientale, Via Cadorna 7, 17100 Savona, Italy

²ISAC-CNR, Istituto di Scienze dell'Atmosfera e del Clima, Corso Fiume 4, 10133 Torino, Italy

Received: 2 November 2007 – Accepted: 6 November 2007 – Published: 22 November 2007

Correspondence to: M. Baudena (mara.b@cima.unige.it)

HESSD

4, 4241–4264, 2007

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏮

⏭

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

We discuss a simplified, conceptual model for the dynamics of the soil-vegetation system in drylands. The model considers the different dynamical processes taking place in vegetated and non-vegetated soil and it distinguishes between the upper soil layer, where rapid evaporation dominates, and the deeper root layer where only plant transpiration takes place. We explore the role of rainfall intermittency and of different plant colonization strategies, and discuss in detail the effect of two different vegetation feedbacks: reduced evaporation due to plant shading and increased infiltration in vegetated areas. The results of the analysis indicate that both temporal rainfall intermittency and the shading/infiltration feedbacks have a beneficial effect on vegetation. However, it turns out that in this model rainfall intermittency and vegetation feedbacks have almost a mutually exclusive role: whenever one of these two components is present, the addition of the other does not further affect vegetation dynamics in a significant way.

1 Introduction

Understanding the interplay of climate and biosphere is one of the intriguing open issues in Earth System Science. In terrestrial environments, vegetation is affected by atmospheric and soil conditions, and at the same time it influences the atmosphere and the soil, through fluxes of water, energy, and carbon dioxide. These interactions are linked with several feedback mechanisms, whose understanding is necessary for a correct description of the global climate system: Accurate climate modeling needs to face the challenge of considering ecosystem dynamics not only as a lower boundary condition to atmospheric flows, but rather as one of the leading actors at play.

The network of feedbacks between vegetation, climate and soil is, in general, difficult to disentangle. For this reason, simplified, conceptual models of climate-vegetation interactions can be of some value to elucidate the basic mechanisms at work and identify relevant parameters and processes. In this work we follow a mechanistic, minimalis-

HESSD

4, 4241–4264, 2007

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tic approach and discuss a simplified model of vegetation-soil interactions in arid and semi-arid regions. We assume that the only limiting factor is water and we make the simplifying hypothesis that plant dynamics depends only on soil moisture, similarly to the approach described in the book of [Rodriguez-Iturbe and Porporato \(2005\)](#). Vegetation dynamics is described by an implicit-space approach, as discussed in the works of [Levins \(1969\)](#) and [Tilman \(1994\)](#). In this model, rainfall is the external input, either kept constant or assumed to be stochastic and intermittent in time.

In a previous work ([Baudena et al., 2007](#), BA in the following), we introduced an eco-hydrological box model, simpler than the one described here, to estimate the effect of intermittent water availability on vegetation dynamics. The results of that work showed that temporal rainfall intermittency allows for vegetation persistence at low values of annual rainfall volume, where it would go extinct if rainfall were constant. Rainfall intermittency also generates long-term fluctuations in vegetation cover, even in the absence of significant inter-annual variations in the statistical properties of precipitation.

On the other hand, vegetation modifies the environment in which it lives in many different ways, other than those addressed in BA. In arid climates, vegetation diminishes evaporation due to its shadowing effects ([Zeng et al., 2006](#); [D’Odorico et al., 2005](#)), and it facilitates infiltration with respect to bare soil surfaces ([Rietkerk and van de Koppel, 1997](#); [West, 1990](#); [Walker et al., 1981](#)), a process that has been extensively analyzed with explicit-space models of vegetation patterns in drylands (e.g. [von Hardenberg et al., 2001](#); [Rietkerk et al., 2002](#); [Gilad et al., 2004, 2007](#)). Both mechanisms result in a positive feedback that favors vegetation persistence.

The simple approach of BA cannot properly capture the two processes described above: In that model, it is not possible to represent the higher infiltration occurring in vegetated soil as compared to bare soil, as soil moisture is spatially averaged over vegetated and non vegetated surfaces. In addition, in BA evaporation was assumed to occur from the whole root layer, whereas in arid and semi-arid regions only the first 5–10 cm of bare soil dry out, owing to the strong and rapid evaporation, while the underlying soil layer remains wet and functions as water storage ([Mahrt and Pan, 1984](#);

**Rainfall intermittency
and vegetation
feedbacks in
drylands**

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Allen et al., 2005; D’Odorico et al., 2005).

To model the infiltration and shading feedbacks, in this work we separately consider the humidity of bare and vegetated soils. Besides, the soil is now divided in two horizontal layers: a thin layer at the surface, where evaporation is intense, and a deeper layer for the underneath root zone. In this way, the model has a total of three soil compartments: the surface layer in either vegetated or non-vegetated soil, and the deeper root layer in vegetated soil. This approach easily allows for a representation of evaporation from the top soil layer only and for a distinction between moisture in vegetated and bare soils.

The rest of this paper is organized as follows. Section 2 provides an introduction to the model adopted here. Section 3 reports the main results of our numerical exploration, and Sect. 4 provides a discussion of our results. Summary and conclusions are reported in Sect. 5.

2 Model description

The model adopted here describes the coupled dynamics of vegetation and soil moisture, and it is an extension of the model discussed in BA. Soil moisture dynamics is described following the approach of Laio et al. (2001), and vegetation dynamics is described by an equation analogous to that introduced by Levins (1969) and Tilman (1994).

The soil is divided into top and bottom layers, allowing the representation of evaporation processes taking place only from the upper layer of soil, whereas from the deeper soil layer only transpiration takes place, due to root uptake. Soil moisture is modeled separately for vegetated and bare soil regions, and differential infiltration occurs between bare soil and vegetation. The system dynamics is modeled by four ordinary differential equations (ODEs) that determine the evolution of four prognostic variables: s_e , the soil moisture in the top layer of bare (empty) soil, Eq. (1); s_u and s_d , the soil moisture in respectively the top and bottom (root) layers of vegetated sites, Eq. (2)–(3);

HESSD

4, 4241–4264, 2007

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

b , the fraction of soil covered by vegetation, Eq. (4):

$$nZ_u \frac{ds_e}{dt} = I(s_e, fr) - E(s_e) - L(s_e) \quad (1)$$

$$nZ_u \frac{ds_u}{dt} = I \left[s_u, r + (1-f)r \frac{1-b}{b+\epsilon} \right] - \sigma E(s_u) - L(s_u) \quad (2)$$

$$nZ_d \frac{ds_d}{dt} = I_d \left[s_u, s_d, r + (1-f)r \frac{1-b}{b+\epsilon} \right] + L(s_u) - T(s_d) - L(s_d) \quad (3)$$

$$\frac{db}{dt} = g(s_e, s_d)b(1-b) - \mu(s_d)b \quad (4)$$

The relative soil moisture content ($0 \leq s_e, s_u, s_d \leq 1$) is averaged over the soil layer (see also e.g. [Laio et al., 2001](#)), n is soil porosity, and Z_u and Z_d are the top and bottom layer depths. The quantity I measures the infiltration rate from daily rainfall r , E is the evaporation rate, T is the transpiration rate, and L represents the water loss due to leakage. Note that we do not explicitly model soil moisture in the deep layer in bare soil, as we assumed that the water stored there cannot be used by plants and it is effectively lost from the system.

Soil moisture dynamics in non-vegetated soil is modeled by Eq. (1). The input is the infiltration from rainfall, I , the output is the sum of evaporation, E , and leakage, L . Rainfall infiltrates in the top soil layer till it saturates, as in [Laio et al., 2001](#):

$$I(s_e, \tilde{r}) = \begin{cases} \tilde{r} & \text{if } \tilde{r} \Delta t < (1-s_e)nZ_u \\ \frac{1-s_e}{\Delta t} nZ_u & \text{if } \tilde{r} \geq \frac{1-s_e}{\Delta t} nZ_u \end{cases} \quad (5)$$

where $\Delta t = 0.1$ days and $\tilde{r} = fr$.

In bare soil, only a fraction f of the total rainfall actually infiltrates, with $0 \leq f \leq 1$. This represents the joint effect of the presence of biophysical crusts that limit infiltration in non-vegetated areas (West, 1990), and of the increased infiltration in vegetated areas due to the presence of roots (Walker et al., 1981; Tietjen et al., 2007¹). Losses in bare soil are due to evaporation and leakage. Since the soil is not vegetated, no transpiration occurs. The evaporation rate is zero below a soil moisture threshold corresponding to the hygroscopic point (s_h), then it increases linearly up to the soil field capacity (s_{fc}), above which it assumes a constant value (BA; Borgogno et al., 2007),

$$E = \begin{cases} 0 & \text{if } s \leq s_h \\ E_w \frac{s - s_h}{s_w - s_h} & \text{if } s_h < s \leq s_{fc} \\ E_w \frac{s_{fc} - s_h}{s_w - s_h} & \text{if } s_{fc} < s \leq 1 \end{cases} \quad (6)$$

E_w is fixed at 0.4 mm per day, so that the maximum evaporation at field capacity, $E(s_{fc})$, is about 3 mm per day, in analogy to e.g. Borgogno et al. (2007); D'Odorico et al. (2005). In BA, E_w was fixed at a lower value because evaporation was assumed to occur from the whole root layer.

Leakage losses occur above the soil field capacity and they are modeled by an exponential growth to the saturated hydraulic conductivity K_s , reached when the soil is

¹Tietjen, B., Zehe, E., and Jeltsch, F.: Modelling water availability to plants in drylands under climate change, Water Resour. Res., under revision, 2007.

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

saturated ($s_e=1$),

$$L(s) = \begin{cases} 0 & \text{if } s \leq s_{fc} \\ K_s \frac{e^{\beta(s-s_{fc})} - 1}{e^{\beta(1-s_{fc})} - 1} & \text{if } s_{fc} < s \leq 1 \end{cases} \quad (7)$$

Equations (2) and (3) represent the dynamics of relative soil moisture content in the vegetated part. Infiltration occurs in the upper layer up to saturating it, as in bare soil, Eq. (5). The excess water that does not infiltrate in bare soil, $(1-f)r$, is assumed to spread rapidly over the vegetated part of the box, and there it participates in the infiltration process together with the water from local rainfall. Referring to the infiltration as expressed by Eq. (5), the water infiltrating in vegetated soil is therefore

$$r' = r + (1-f)r \frac{1-b}{b+\epsilon}, \quad (8)$$

where $\frac{1-b}{b+\epsilon}$ is a weight used to distribute over the vegetated soil the water which is not infiltrating in bare, crusted areas. To avoid unphysical divergence of infiltration when $b \rightarrow 0$, we introduce a (small) regularizing constant, $\epsilon=0.01$.

When the top layer saturates, water starts infiltrating into the bottom layer, in analogy to the one-layer approach of e.g. [Rodriguez-Iturbe and Porporato \(2005\)](#). Infiltration I_d

into the bottom layer is thus simply modeled as:

$$I_d(s_u, s_d, r') = \begin{cases} 0 & \text{if } r' \Delta t < (1 - s_u)nZ_u \\ r' - \frac{1 - s_u}{\Delta t} nZ_u & \text{if } r' \Delta t - (1 - s_u)nZ_u + \\ & + L(s_u)\Delta t < (1 - s_d)nZ_d \\ \frac{1 - s_d}{\Delta t} nZ_d - L(s_u) & \text{if } r' \Delta t - (1 - s_u)nZ_u + \\ & + L(s_u)\Delta t \geq (1 - s_d)nZ_d \end{cases} \quad (9)$$

where $\Delta t = 0.1$ days. If the bottom layer gets saturated, water is lost to still deeper layers and cannot be directly used by plant roots. This representation of infiltration is strongly simplified (for a detailed representation, see e.g. [Rigby and Porporato, 2006](#)).

5 Evaporative losses E occur only from the top layer, with the same dependence on soil moisture content expressed by Eq. (6) for bare soil. Vegetation exerts a shadowing effect on the soil, limiting evaporation. This effect is included in the parameter σ , which multiplies E and can vary from 0 to 1, representing respectively a total or a null shadowing effect from vegetation cover.

Transpiration contributions from the top layer are implicitly included in the evaporation term, because transpirational losses from the first few centimeters of soil are negligible compared to vegetation shadowing effects ([Scanlon and Albertson, 2003](#); [Borgogno et al., 2007](#); [D'Odorico et al., 2005](#)). Thus, we explicitly include a transpiration term T lifting water into the atmosphere from the bottom layer only, as a function of its soil moisture content, s_d . The transpiration rate is set to zero below the wilting point, s_w ; it grows linearly above s_w and assumes a constant value E_{\max} at s^* , following

e.g. [Rodriguez-Iturbe et al., 1999](#):

$$T(s) = \begin{cases} 0 & \text{if } s \leq s_w \\ E_{\max} \frac{s - s_w}{s^* - s_w} & \text{if } s_w < s \leq s^* \\ E_{\max} & \text{if } s \geq s^* \end{cases} \quad (10)$$

Leakage losses from the bottom of the soil layers are represented as in Eq. (7). Water percolating from the top layer, $L(s_u)$ enter the bottom layer. This phenomenon is different from the infiltration because it happens on slower time scales, due to gravity and not to root channeling.

In our approach, vegetation is modeled in Eq. (4) by an implicit-space logistic equation for the fraction of space, b , occupied by plants ([Tilman, 1994](#)). The pro-capite colonization rate, g , represents the ability of plants to colonize bare sites, while sites become empty due to the local extinction rate, μ . The pro-capite colonization rate multiplies the fraction of space covered by vegetation, b , and the fraction of empty space, $1-b$, to express the total growth rate of the fraction of vegetated sites. Therefore, g represents both the plant ability to produce seeds and the germination probability of the seeds. As in BA, the colonization rate depends on soil moisture. In principle, the colonization rate depends on the soil moisture of both empty and vegetated soil, and it is factorized in a seed production term, g_b , and a seed germination term, g_e :

$$g = g_0 g_e(s_e) g_b(s_b) . \quad (11)$$

- 5 Here, g_0 is the maximum colonization rate, $g_e(s_e)$ represents the seed establishment probability as a function of surface humidity in bare soil, and $g_b(s_b)$ represents the plant ability to produce seeds as a function of soil moisture in vegetated sites.

The seed production term, g_b , is assumed to vary with the soil moisture of the vegetated part, s_b . For simplicity, in the following we choose $s_b = s_d$. The results

HESSD

4, 4241–4264, 2007

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

do not show qualitative changes when s_b is chosen differently, e.g. equal to the weighted average of the moisture of the top and bottom soil layers in vegetated soil, $s_b = (s_u Z_u + s_d Z_d) / (Z_u + Z_d)$. The factor g_b is supposed to be zero below a threshold corresponding to the wilting point, s_w , to grow linearly with soil moisture below the threshold for fully open stomata, s^* , and to become constant and equal to one for $s_d > s^*$:

$$g_b = \begin{cases} 0 & \text{if } s_d < s_w \\ \frac{s_d - s_w}{s^* - s_w} & \text{if } s_w \leq s_d < s^* \\ 1 & \text{if } s^* \leq s_d \end{cases} \quad (12)$$

The factor g_e represents the fact that seeds can germinate only when the humidity of the bare soil where they lie is sufficiently large. Dependence of seed germination on moisture of bare soil is modeled by assuming that g_e varies from 0 to 1 with a hyperbolic tangent shape centered around a threshold value, chosen as the plant wilting point, s_w , analogously to the approach of BA:

$$g_e(s_e) = \frac{1}{2} \left[1 + \tanh \frac{s_e - s_w}{a} \right] \quad (13)$$

The mortality rate is assumed to depend on soil moisture where plants grow. In particular, μ depends on the bottom soil moisture s_d : μ tends to the value μ_1 for $s < s_w$ and to the value μ_2 for $s > s_w$, where $\mu_1 > \mu_2$. Following BA, we assume a hyperbolic tangent shape centered around s_w ,

$$\mu(s_d) = \frac{\mu_1 + \mu_2}{2} + \frac{\mu_1 - \mu_2}{2} \tanh \frac{s_w - s_d}{a} \quad (14)$$

For illustration, Fig. (1) shows $g_0 g_b$ and μ as a function of s_d and $g_0 g_e$ as a function of s_e .

The values of the model parameters used in the following are reported in Tab. 1, and we have assumed to deal with annual vegetation on a loamy soil. Other choices of soil properties lead to qualitatively analogous results.

3 Results

In this section we report the results of our numerical exploration, focussing on the effect of vegetation feedbacks (infiltration and shading), and comparing the response of vegetation with different colonization strategies. We also estimate the impact of different rainfall regimes: either a constant input or temporally intermittent rainfall forcing. As in Laio et al. (2001) and in BA, precipitation intermittency is obtained by assuming independent, instantaneous rainfall events distributed as a Poisson process with interarrival time N . The intensity of each event is extracted from an exponential distribution with average rainfall depth r_0 . Rainfall occurs only during the wet season (200 days), and the soil-vegetation system is assumed to be “frozen” during the dry season, see the discussion on this issue in BA.

3.1 Colonization rate dependent only on soil moisture in the root layer.

First we consider a case where the colonization rate, g , depends only on the vegetation ability to produce propagules, and not on the ability of seeds to germinate in bare soil. Consistently, we assume

$$g(s_b)=g_0g_b(s_b), \tag{15}$$

with g_b defined as in Eq. (12) and $g_e(s_e) \equiv 1$.

Figure 2a shows the vegetation behavior and Fig. 3a shows the soil moisture content when no vegetation feedbacks are introduced in the system: The bare soil is not crusted ($f=1$) and vegetation does not decrease evaporation ($\sigma=1$). In case of constant rainfall, vegetation is not present below about 800 mm per year. As in the case

considered by BA, rainfall intermittency favors vegetation, shifting to about 400 mm per year the threshold for vegetation persistence. The surface soil moisture s_e and s_u are identical because no feedback is present in this case. In the intermittent case, surface soil moisture is lower than in the constant case, due to increased leakage and deep infiltration, which increase soil moisture in the root layer, s_d .

Figures 2b and 3b show the results when only a shading feedback is introduced, with $\sigma=0.5$. The equilibrium solution for constant rainfall is significantly affected by the shading feedback, with vegetation appearing for less than 500 mm per year. The moisture content of bare soil is not affected, but the soil moisture in the top vegetated layer, s_u , and, consequently, s_d , increase (Fig. 3b). Vegetation is again favored by intermittent rainfall but the advantage is lower than for the case with no feedbacks. Soil moisture in the bottom soil layer is higher under intermittent than under constant rainfall, but the difference is smaller than in the case without shading feedback.

Figures 2c and 3c show the results for the case where only the infiltration feedback is present: here, the biogenic crust is assumed to limit infiltration in bare soil to 70% of total precipitation ($f=0.7$). Direct estimates in the Negev desert indicate that only about 60–80% of the precipitated water infiltrates over bare soil (M. Shachak, personal communication). Run-off water from crusted, bare soil infiltrates in vegetated patches (here, we assume that no run-off water is lost due to evaporation). This mechanism leads to vegetation presence at very low rainfall, both in the intermittent and in the constant precipitation cases, thanks to an increase in water availability in the vegetated parts (higher s_u and s_d). This increase is accompanied by a decreased value of s_e , see Fig. 3c.

When the two feedbacks are considered together, as shown in Figs. 2d and 3d, the combined effect is a definite advantage for vegetation. Surprisingly, this advantage is more pronounced in the case of constant rainfall than for intermittent precipitation. In the constant case, vegetation is present even at values of total annual rainfall lower than 100 mm per year. The soil moisture in the root layer, s_d , is only slightly larger for intermittent rainfall than for constant rainfall. Under these conditions, rainfall intermit-

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tency plays a minor role: apparently, the presence of vegetation feedbacks reduces the role of rainfall intermittency.

3.2 Colonization rate dependent on soil moisture in the root layer and in bare soil.

A different approach consists in modeling vegetation colonization rate as dependent on both seed establishment probability and plant status. Seeds germinate over bare soil, and in this version of the model we assume that the colonization is expressed as

$$g=g_0g_e(s_e)g_b(s_b)$$
 (16)

with g_e as in (13) and g_b as in (12).

5 The case without feedbacks, shown in Figs. 4a and 5a, is identical to the case without dependence on bare soil moisture (Fig. 2a, because bare soil limit vegetation growth only at low water amounts, for $s_e < s_w$, where in any case $g_b \equiv 0$, owing to the fact that $s_d \leq s_w$, and thus vegetation cover is zero. Also in the cases with a presence of vegetation feedbacks the results are very similar to those obtained without dependence
10 on bare soil humidity ($g=g_0g_b$). In particular, Fig. 5a and 5b are identical to Fig. 3a and 3b. Indeed, soil moisture dynamics changes with vegetation cover only when the differential infiltration between vegetated and bare surface is included (Figs. 3c,d).

The effect of the dependence of the colonization rate on s_e manifests itself in two features. First, under constant rainfall conditions, a limit is imposed to vegetation growth
15 when s_e is lower than s_w . As shown in Figs. 4c and 4d, $b > 0$ only above a value of annual rainfall of about 100 mm per year. Second, the beneficial effect of rainfall intermittency over vegetation is smaller, and it almost disappears when both feedbacks are considered.

4 Discussion

20 The results illustrated in the previous section suggest a number of considerations on the role of vegetation feedbacks, on rainfall intermittency and on their interplay.

Rainfall intermittency
and vegetation
feedbacks in
drylands

M. Baudena et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏮	⏭
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

When g depends only on s_d , the minimum constant rainfall needed to guarantee vegetation survival is ≈ 800 mm per year, corresponding to the threshold at which s_d grows above s_w , and thus $g > 0$. In the absence of any vegetation feedbacks, the non-linear dependence of g on s_d leads to a significant beneficial effect of rainfall intermittency, similarly to what has been discussed in detail by BA. The situation is analogous when g depends on both s_e and s_d .

In the presence of vegetation feedbacks, the situation can change. Vegetation limits surface evaporation, increasing water in the top layer of vegetated soil. Deep infiltration and leakage grow, thus increasing deep water storage and advantaging vegetation growth owing to the dependence of g on s_d . Interestingly, Figs. 2b and 4b show that the larger advantage due to shading occurs when rainfall is homogeneously distributed in time.

The other effect of vegetation is to facilitate infiltration from rainfall, as compared to bare soil where crusts of either physical or biological origin limit soil infiltrability (West, 1990; Rietkerk et al., 2002; Gilad et al., 2004). This effect is always favorable to vegetation because it increases biomass locally. On the other hand, it is more difficult for seeds to germinate over dryer soils. When g depends only on s_d , infiltration feedback has a strong positive effect on b , which remains positive, albeit small, even at very low annual rainfall. When g depends also on s_e , for a constant precipitation input vegetation appears at slightly larger values of annual rainfall, but still much smaller than for the case without the infiltration feedback.

The infiltration feedback is considered to be the responsible for vegetation pattern formation in spatially explicit models, bringing advantages to vegetation (von Hardenberg et al., 2001; Rietkerk et al., 2004). In these models, biomass grows locally due to the larger amount of water available for root uptake. One can include this effect by considering higher colonization ability when the soil in the root layer is wetter, representing the fact that plants can colonize more easily sites around them. Introducing this dependency, infiltration and shading feedbacks have larger effects, in agreement with the results obtained with explicit-space models.

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

For the case with the two vegetation feedbacks acting together, the advantage due to the feedbacks is larger in case of constant rainfall than for intermittent rainfall. Both feedbacks increase surface moisture in the vegetated parts and therefore deep water storage. A similar effect is obtained in the case of intermittent rainfall occurring in high-amplitude intermittent pulses, but soil moisture cannot grow indefinitely, and the sum of the two effects (rainfall intermittency and vegetation feedbacks) is practically not larger than the effect of each of them. In general, we also observe that the feedbacks are more advantageous, in both rainfall regimes, when g does not depend on bare soil moisture, which is significantly lowered by the infiltration feedback.

In the exploration discussed above, we did not explicitly consider the case when the colonization rate, g , depends only on soil moisture in bare soil, i.e.,

$$g(s_e) = g_0 g_e(s_e) \quad (17)$$

where $g_e(s_e)$ varies as in Eq. (13), with a threshold around $s_e = s_w$. Such a choice could be motivated by the fact that in arid lands, the existence of a *seed bank* is often considered as a plausible hypothesis. Seeds are present everywhere and ready to germinate under favorable conditions, due to temporal or spatial variability (e.g. Oren, 2001). Under such conditions, dependence of colonization ability on just the soil moisture in empty sites is a possible modeling choice, and moisture in the root layer comes into play only in plant mortality. Model runs with this choice for g indicate that in this case the infiltration and shading feedbacks favor vegetation only slightly, through diminished mortality, while spread in new sites is limited because of increased difficulty in colonizing bare soil. In fact, in this case the infiltration feedback could even reduce the spatial spread of vegetation, favoring instead the growth of new vegetation in patches already occupied by other older plants.

We believe that an interesting point raised by the results discussed above is that the influence of vegetation feedbacks is larger when rainfall is kept constant in time. Many studies dealing with the importance of vegetation feedbacks, both with implicit and explicit spatial structure, considered rainfall as a constant input (e.g. Zeng et al.,

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2006; von Hardenberg et al., 2001; Rietkerk et al., 2002). Here the importance of rainfall intermittency is underlined again, showing that part of the effects due to the feedbacks (namely, deep water storage) can be obtained, without feedbacks, when rainfall occurs in intermittent pulses. Depending on the vegetation strategy and on the type of vegetation feedbacks present in the system, rainfall intermittency can thus be more or less beneficial to vegetation.

5 Summary and conclusion

In this work we have discussed a simple vegetation-soil moisture model for drylands. The model includes two effects which were discarded in most previous models: Evaporation occurs from the top layer only, allowing water storage in the deeper root layer, and vegetation feedbacks lead to reduced evaporation from vegetated areas and to differential infiltration between vegetated and empty sites.

Plant colonization behavior has been assumed to follow different strategies, depending on soil moisture of the bare soil, to represent seed germination abilities, and on the moisture in vegetated soil, to represent seed production. Depending on the functional form of the colonization rate, the effect of infiltration or shading feedbacks is different. In general, however, we find that in this model the beneficial effects of rainfall intermittency and of vegetation feedbacks are somehow mutually exclusive: vegetation feedbacks act at their maximum strength for a constant rainfall input, and rainfall intermittency loses most of its beneficial power when strong vegetation feedbacks are present. One can thus expect that plants living in areas with different rainfall climatologies (more or less intermittent for the same value of rainfall volume) may have evolved different strategies.

Acknowledgements. We are grateful to F. D’Andrea, S. Dekker, J. von Hardenberg, A. Porporato, M. Rietkerk and M. Shachak for useful discussions and comments on this work.

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

References

- Allen, R. G., Pereira, L. S., Smith, M., Raes, D., and Wright, J. L.: FAO-56 Dual Crop Coefficient Method for Estimating Evaporation from Soil and Application Extensions, *J. Irrig. Drain. E.*, 131, 2–13, 2005. [4244](#)
- 5 Baudena, M., Boni, G., Ferraris, L., von Hardenberg, J., and Provenzale, A.: Vegetation response to rainfall intermittency in drylands: Results from a simple ecohydrological box model, *Adv. Water Res.*, 30, 1320–1328, 2007. [4243](#)
- Borgogno, F., D'Odorico, P., Laio, F., and Ridolfi, L.: Effect of rainfall interannual variability on the stability and resilience of dryland plant ecosystems, *Water Resour. Res.*, 43, W06411, doi:10.1029/2006WR005314, 2007. [4246](#), [4248](#)
- 10 D'Odorico, P., Laio, F., and Ridolfi, L.: Noise-induced stability in dryland plant ecosystems, *P. Natl. A. Sci.*, 102, 10 819–10 822, 2005. [4243](#), [4244](#), [4246](#), [4248](#)
- Gilad, E., von Hardenberg, J., Provenzale, A., Shachak, M., and Meron, E.: Ecosystem Engineers, in : from Pattern Formation to Habitat Creation, *Phys. Rev. Lett.*, 93, 098105, 2004. [4243](#), [4254](#)
- 15 Gilad, E., von Hardenberg, J., Provenzale, A., Shachak, M., and Meron, E.: A mathematical model of plants as ecosystem engineers, *J. Theor. Biol.*, 244, 680–691, 2007. [4243](#)
- Laio, F., Porporato, A., Ridolfi, L., and Rodriguez-Iturbe, I.: Plants in water controlled ecosystems, in: active role in hydrological processes and response to water stress II. Probabilistic soil moisture dynamics, *Adv. Water Res.*, 24, 707–723, 2001. [4244](#), [4245](#), [4251](#)
- 20 Levins, R.: Some demographic and genetic consequences of environmental heterogeneity for biological control, *Bulletin of the Entomological Society of America*, 15, 237–240, 1969. [4243](#), [4244](#)
- Mahrt, L. and Pan, H.: A two layer model of soil hydrology, *Bound.-Lay. Meteorol.*, 29, 1–20, 1984. [4243](#)
- 25 Oren, Y.: Patchiness, disturbances, and the flows of matter and organisms in an arid landscape, in: a multiscale experimental approach, Ph.D. thesis, Ben-Gurion University of the Negev, Beer-Sheva, Israel, 2001. [4255](#)
- Rietkerk, M. and van de Koppel, J.: Alternate stable states and threshold effects in semi-arid grazing systems, *Oikos*, 79, 69–76, 1997. [4243](#)
- 30 Rietkerk, M., Boerlijst, M. C., van Langevelde, F., HilleRisLambers, R., van de Koppel, J., Kumar, L., Prins, H. H. T., and Roos, A. M. D.: Self-Organization of Vegetation in Arid Ecosys-

HESSD

4, 4241–4264, 2007

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- tems, *American Naturalist*, 160, 524–530, 2002. [4243](#), [4254](#), [4256](#)
- Rietkerk, M., Dekker, S. C., de Ruiter, P. C., and van de Koppel, J.: Self-Organized Patchiness and Catastrophic Shifts in Ecosystems, *Science*, 305, 1926–1929, 2004. [4254](#)
- Rigby, J. and Porporato, A.: Simplified stochastic soil moisture models: a look at infiltration, *Hydrol. Earth Syst. Sci. Discuss.*, 3, 1339–1367, 2006,
<http://www.hydrol-earth-syst-sci-discuss.net/3/1339/2006/>. [4248](#)
- Rodriguez-Iturbe, I. and Porporato, A.: *Ecohydrology of water controlled ecosystems: soil moisture and plant dynamics*, Cambridge University Press, doi:10.2277/0521819431, 2005. [4243](#), [4247](#)
- Rodriguez-Iturbe, I., D’Odorico, P., Porporato, A., and Ridolfi, L.: On the spatial and temporal links between vegetation, climate and soil moisture, *Water Resour. Res.*, 35, 3709–3722, 1999. [4249](#)
- Scanlon, T. M. and Albertson, J. D.: Inferred controls on tree/grass composition in a savanna ecosystem, in: *Combining 16-year normalized difference vegetation index data with a dynamic soil moisture model*, *Water Resour. Res.*, 39, 1224, doi:10.1029/2002WR001881, 2003. [4248](#)
- Tilman, D.: Competition and biodiversity in spatially-structured habitats, *Ecology*, 75, 2–16, 1994. [4243](#), [4244](#), [4249](#)
- von Hardenberg, J., Meron, E., Shachak, M., and Zarmi, Y.: Diversity of Vegetation Patterns and Desertification., *Phys. Rev. Lett.*, 87, 198 101, 2001. [4243](#), [4254](#), [4256](#)
- Walker, B. H., Ludwig, D., Holling, C. S., and Peterman, R. M.: Stability of semi-arid savanna grazing system, *J. Ecol.*, 69, 473–498, 1981. [4243](#), [4246](#)
- West, N. E.: Structure and function of soil microphytic crusts in wildland ecosystems of arid to semi-arid regions, *Adv. Ecol. Res.*, 20, 179–223, 1990. [4243](#), [4246](#), [4254](#)
- Zeng, X., Wang, A., Zeng, Q., Dickinson, R. E., Zeng, X., and Shen, S. S. P.: Intermediately complex models for the hydrological interactions in the atmosphere-vegetation-soil system, *Adv. Atmos. Sci.*, 23, 127–140, 2006. [4243](#), [4255](#)

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Rainfall intermittency
and vegetation
feedbacks in
drylands**

M. Baudena et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. List of model parameters and of their values. Units are expressed in millimeters (mm) and days (d). The chosen values are appropriate for annual vegetation (grasses) on loamy soil.

Symbol	Meaning	Value	Units
n	Soil porosity	0.45	
Z_u	Depth of the upper soil layer	50	mm
Z_d	Depth of the deeper soil layer	250	mm
s_h	Soil hygroscopic point	0.19	
s_w	Soil wilting point	0.24	
s^*	Soil moisture value at which plants start closing their stomata	0.57	
s_{fc}	Soil field capacity	0.65	
E_{max}	Maximum evapotranspiration at s^*	4.5	mm d ⁻¹
E_w	Potential evapotranspiration at wilting point	0.4	mm d ⁻¹
K_s	Saturated hydraulic conductivity	200	mm d ⁻¹
r_0	Average rainfall depth	15	mm d ⁻¹
N	Average rainfall interarrival time	5	d
β	Water retention parameter	14.8	
g_0	Vegetation maximum colonization rate	2.2×10^{-3}	d ⁻¹
μ_2	Vegetation maximum extinction rate	0.3×10^{-3}	d ⁻¹
μ_1	Vegetation minimum extinction rate	2.7×10^{-3}	d ⁻¹
a	Hyperbolic tangent width	0.002	
f	Fraction of water infiltrating over bare soil	0.7	
σ	Shading effect due to vegetation	0.5	

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.




Fig. 1. Left panel: Continuous line, colonization rate depending on soil moisture in the root layer, s_d , $g = g_0 g_b(s_d)$. Dotted line, mortality rate as a function of s_d . Right panel: Colonization rate depending on soil moisture of bare sites, $g = g_0 g_e(s_e)$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.




Fig. 2. Vegetation cover, b , versus annual rainfall, for the case where the colonization rate depends only on the soil moisture of the root layer (s_d). Continuous lines represent equilibrium points in the case of constant rainfall, crosses represent average vegetation cover in the case of intermittent rainfall. The four panels refer to the following situations: **(a)** No vegetation feedbacks; **(b)** Presence of shading feedback only; **(c)** Presence of infiltration feedback only; **(d)** Presence of both infiltration and shading feedbacks. Rainfall is expressed by its annual value, i.e. the total rainfall during the wet season. The total annual rainfall is varied by changing the average interarrival time. Parameter values are reported in Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.




Fig. 3. Soil moisture, s , of bare and vegetated soil versus annual rainfall, for the case where the colonization rate depends only on the soil moisture of the root layer (s_d). Lines represent equilibrium points in the case of constant rainfall (Continuous line, s_u , dotted line, s_d , dash-dotted line, s_e). Symbols represent the average vegetation cover for intermittent rainfall: + marks for s_u , circles for s_d , x-signs for s_e . The values of s_u and s_d are not plotted when $b \rightarrow 0$. The four panels refer to the following situations: **(a)** No vegetation feedbacks; **(b)** Presence of shading feedback only; **(c)** Presence of infiltration feedback only; **(d)** Presence of both infiltration and shading feedbacks. Rainfall is expressed by its annual value, i.e. the total rainfall during the wet season. The total annual rainfall is varied by changing the average interarrival time. For parameter values see Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.




Fig. 4. Vegetation cover, b , versus annual rainfall, for the case where the colonization rate depends on both the soil moisture of the root layer in vegetated soil and on that of bare soil. Continuous lines represent equilibrium points in the case of constant rainfall, crosses represent the average vegetation cover in the case of intermittent rainfall. The four panels refer to the following situations: **(a)** No vegetation feedbacks; **(b)** Presence of shading feedback only; **(c)** Presence of infiltration feedback only; **(d)** Presence of both infiltration and shading feedbacks. Rainfall is expressed by its annual value, i.e. the rainfall during the wet season. The total annual rainfall is varied by changing the average interarrival time. Parameter values are reported in Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Rainfall intermittency and vegetation feedbacks in drylands

M. Baudena et al.




Fig. 5. Soil moisture, s , of bare and vegetated soil versus annual rainfall, for the case where the colonization rate depends on both the soil moisture of the root layer in vegetated soil and on that of bare soil. Lines represent equilibrium points in the case of constant rainfall (Continuous line, s_u , dotted line, s_d , dash-dotted line, s_e), symbols represents average vegetation cover in the case of intermittent rainfall (+marks for s_u , circles for s_d , x-signs for s_e). The values of s_u and s_d are not plotted when $b \rightarrow 0$. The four panels refer to the following situations: **(a)** No vegetation feedbacks; **(b)** Presence of shading feedback only; **(c)** Presence of infiltration feedback only; **(d)** Presence of both infiltration and shading feedbacks. Rainfall is expressed by its annual value, i.e. the rainfall during the wet season. The total annual rainfall is varied by changing the average interarrival time. For parameter values see Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion